



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Respiration in wood: integrating across tissues, functions and scales

Citation for published version:

Meir, P, Mencuccini, M & Coughlin, SI 2020, 'Respiration in wood: integrating across tissues, functions and scales', *New Phytologist*, vol. 225, no. 5, pp. 1824-1827. <https://doi.org/10.1111/nph.v225.5>

Digital Object Identifier (DOI):

[10.1111/nph.v225.5](https://doi.org/10.1111/nph.v225.5)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

New Phytologist

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Respiration in wood: integrating across tissues, functions and scales

Commentary on Salomón et al. (2019), 'TReSpire – a biophysical TRee Stem respiration model', *New Phytologist*.

~~Patrick-Patrick~~ Meir^{1,2*}, ~~Maurizio-Maurizio~~ Mencuccini^{3,4} & ~~S~~ Ingrid Coughlin¹

1. Research School of Biology, Australian National University, Canberra ACT 2601, Australia.
2. School of Geosciences, University of Edinburgh, Edinburgh EH9 3 FF, UK.
3. ICREA, Pg. Lluís Companys 23, 08010 Barcelona (Spain).
4. CREA, Universidad Autonoma de Barcelona, Cerdanyola del Valles 08193, Barcelona, Spain

*patrick.meir@anu.edu.au

[+61\(0\)2 6125 8463](tel:+61(0)2 6125 8463)

Word count: 1898

Commented [PM1]: Nichola – Ingrid uses her middle name; her full name is Sarah Ingrid Coughlin. I checked: Ingrid is happy with S. Ingrid Coughlin, but if you need both names, then, the S is Sarah.

Success in life for any seedling requires the simultaneous acquisition of resources from soil and atmosphere. This must be coupled to an ability to distribute these resources to the points of use above and below ground, and to house the distribution system within resilient woody structures. The cells comprising these lignified structures – roots, stems and branches – need to be constructed, maintained and defended. To do so requires chemical energy which is almost entirely derived from respiration. However, despite the fundamental role woody tissue plays in driving life in most terrestrial ecosystems, it has received much less research attention than has the physiology of its greener counterpart, the leaf. To some extent this difference reflects the challenge of understanding the metabolically distinct tissues and related fluid and solute movements that occur simultaneously in wood. Advanced models of xylem and phloem function and of cambial growth have existed for a few years (e.g., Hölttä et al. 2007, 2010, de Schepper et al. & Steppe 2010, Mencuccini et al. 2013), but Salomón et al.'s new paper (NP page details) takes a notable step forward integrating and advancing these approaches. Their model 'TReSpire' combines representations of the essential properties of both phloem and xylem, the use of phloem-derived sugars to produce irreversible cellular growth, together with the activity and physical properties of associated tissues (e.g., cambium, parenchyma, bark). TReSpire describes the variation in the flux of carbon dioxide (CO₂) from tree bark, its underlying respiratory production by both growth and maintenance processes, and its transport, limited by energy expenditure and the fundamental physical constraints of pH, concentration gradients, stem carbon density and diffusive resistances.

Commented [PM2]: Pls note need to add in proofs

The flux of CO₂ from woody tissue is thought to comprise up to 40% of total respiration from vegetation (Carnioli et al. 2016, Yang et al. 2016). Hence, quantifying how it responds to disturbance or climate has substantial consequences across scales, from the carbon budgets of individual plants to the terrestrial biosphere (Meir & Grace 2002, Reich et al. 2006, Huntingford et al. 2017, West et al. 2019). However, woody tissue respiration has proved difficult to quantify accurately. Most efforts have used chamber-based measurements of CO₂ efflux from tree stems (E_a) usually at 1-2 m above ground, then related these fluxes to sap wood volume, or to tree-size metrics (woody tissue area or volume), and assumed a simplified tree structure to extrapolate to whole trees or stands (Ryan et al. 1997, Meir et al. 2017).

Even before scaling to a whole tree, the physiological signals in E_a require interpretation because they reflect multiple sink and source processes occurring below the outer bark. In addition to cellular respiration, E_a fluxes may be affected by CO₂ carried in the transpiration stream (Levy et al. 1999, Teskey et al. 2008), by biochemical consumption, e.g., through cortical photosynthesis or PEP-carboxylase metabolism (Angert et al. 2012), and by diffusion constraints imposed by tissue properties and stem size (Steppe et al. 2007). Salomón et al. have formalised much of this complexity for the first time in a combined model that fuses bark and xylem hydraulics with two additional sub-models that quantify carbohydrate use in

66 respiratory metabolism and growth, together with the dynamics of phloem unloading and
67 consequent carbon balance.

68 The determination of respiration in Salomón et al.'s model is anchored in the classical
69 growth-maintenance paradigm (Thornley 1970, Amthor 2000). This assumes a constant
70 construction respiration cost for biomass of a given elemental composition, and uses live
71 tissue nitrogen content and temperature to drive short-term responses in maintenance
72 respiration (Cannell and Thornley 2000). The simulated mass balance ultimately links
73 fluctuations in cell turgor and stem diameter all the way through to E_a values (Figs 1,2,
74 Salomón et al., this issue). A key advance is that the processes and storage terms
75 determining E_a can be resolved separately at sub-daily temporal resolution as well as over
76 the longer term. The model outputs are broadly consistent with observations reported
77 elsewhere as well as with the detailed experimental measurements made to test the model.
78 This provides initial confidence in the model outputs, despite the sparse validation data for
79 some variables, such as non-structural carbohydrate concentration.

80

81 Besides a more integrated representation of respiratory processes, TReSpire also aims to
82 reduce some of the uncertainties that still surround stem growth. Irreversible radial growth
83 consists of the processes of cambial cell differentiation, cellular periclinal division, radial
84 enlargement, subsequent wall thickening *via* lignification of the secondary wall and final
85 programmed cell death (Rossi et al. 2006). High-precision dendrometry and/or repeated
86 micro-coring coupled with wood anatomical analysis are generally used to quantify these
87 processes. The first method (also employed here) provides rich time-series of highly time-
88 resolved data. However, at least three processes contribute to the observed
89 shrinking/swelling signals measured by dendrometers at the hourly and daily scale, i.e., bark
90 hydraulic capacitance driven by xylem water potential (Zweifel [et al.](#) 2016), radial water
91 transport associated with osmotic changes in bark (de Schepper [et al.](#) & [Steppe](#) 2010) and/or
92 xylem and irreversible growth. Disentangling these processes has become possible recently
93 (Mencuccini et al. 2017) and these new advances are incorporated in TReSpire. By contrast,
94 micro-coring provides less capacity to resolve processes at the daily scale and is inherently
95 destructive by nature. However, compared to dendrometry, it has significant advantages in
96 its capacity to identify individual phenological phases and thus attribute metabolic costs to
97 each one of them at the seasonal time scale (Rossi et al. 2006; Cuny et al., 2015).

98

99 Because interpreting dendrometry signals at diurnal and seasonal time scales remains
100 challenging, fundamental questions over the relative roles of hydraulic (e.g., turgor) versus
101 metabolic (e.g., carbon supply, carbon demand) variables on the control of wood growth
102 have remained unanswered. The insights provided by TReSpire are exploited to separate the
103 detailed time-dynamics not only of respiratory fluxes but also of the subtending irreversible
104 growth processes. Such an approach has the potential to help address outstanding
105 questions, including the impact of high vapour pressure deficit or reduced soil water
106 availability on the rate of radial growth. Coupled with additional measurements of the rate

107 of photosynthesis and carbohydrate content, it may also shed light on the balance between
108 storage, source-driven and sink-driven processes. Interestingly, the authors report that the
109 stem growth rate (in sucrose equivalents) of a young Norway maple tree was lowest in late
110 afternoon, accelerated during the night and peaked in early morning, remaining relatively
111 high also during the middle part of the day. This finding casts doubts over the generally held
112 view that radial xylem growth occurs only during night-time periods of maximum cellular
113 turgor, but is consistent with growth pattern reports for some *Eucalyptus* species
114 (Mencuccini et al. 2017).

115

116 Some of the key measurements used to calibrate the model included high-resolution stem
117 diameter changes, sap flux and chamber-based E_a fluxes. Given the significance of
118 dendrometric measurements for the present analysis and the interest in establishing
119 dendrometric monitoring networks in carbon cycle studies, it is now a priority to examine
120 the comparability of the widely different approaches employed for this purpose.
121 Dendrometer systems vary widely in their design and capacity to account for radial changes
122 in xylem and bark, among other factors. They also differ depending on whether they
123 measure point radius, diameter and/or circumference changes. Their temperature
124 sensitivity and the significance of temperature and water content changes in various tissue
125 compartments (including potentially the heartwood of large trees) will need to be examined
126 carefully.

127

128 A new overall view is thus presented by Salomon et al. of how respiration can vary hourly in
129 the context of stem hydraulic dynamics and growth, with quantitative account also taken of
130 the effects of changes in the transport of CO_2 in the xylem stream and in the changes in
131 gaseous and dissolved CO_2 concentrations in different tissues. For example, detailed *in situ*
132 measurements have shown that E_a fluxes can often be less than the overall respiration rate
133 (R_s) of underlying cells (Angert et al. 2012, Teskey et al. 2008). TReSpire replicates this
134 mechanistically: over a 24 hr cycle, the modelled ratio of E_a/R_s varies between 0.7 and 1,
135 matching field measurements (Angert et al. 2012, Hilman et al. 2018). However, this
136 agreement breaks down at sub-daily timescales, where diurnal E_a/R_s ratios are more stable
137 (Hilman et al. 2018) than the modelled outputs, which overestimate E_a at night. A
138 discrepancy of this sort highlights the advantages accrued by comparing a mechanistic
139 model with field data, and points to the need not just to understand additional processes
140 causing the removal of locally respired CO_2 (Angert et al. 2012), but also when different
141 processes might dominate the signals in E_a and R_s .

142

143 TReSpire has only been validated against saplings thus far, and whilst data on E_a remain
144 relatively sparse for all ecosystems (Yang et al. 2016), it is clear that the relationship
145 between E_a and woody limb size varies strongly. E_a has often been represented as a function
146 of sapwood volume, outer bark diameter and/or bark surface area (e.g., Ryan et al. 1997,
147 Levy et al. 1998, Chambers et al. 2004), though the increased CO_2 transport associated with

148 the larger sap flux that occurs in bigger trees was recently found to explain much variation
149 in E_a with stem size in one species, *Liriodendron tulipifera* (Fan et al. 2017). TReSpire could be
150 used to simulate similar datasets, and in combination with field data, to explore the
151 mechanistic underpinnings of E_a and R_s in large trees under varying environmental
152 conditions (e.g., Metcalfe et al. 2010). Whilst it is unclear how well the classical assumptions
153 of constant cost for different respiration components will hold over large changes in
154 temperature or moisture availability (O’Leary et al. 2019), the integration of hydraulics and
155 respiration in one model may help interpret how respiration varies with limb size and
156 growth conditions.

157
158 To proceed from limb to whole tree, a fine-scale model requires additional information on
159 tree structure. The difficulty of acquiring such data has until now forced a reliance on
160 destructive harvests to enable extrapolation of chamber-based E_a measurements to
161 estimate CO₂ emissions from whole trees. Whilst some of this work has yielded apparently
162 general power-law descriptions of respiration with total biomass (Mori et al. 2010),
163 sensitivity to the power-law exponents is high, and the approach has not yet fitted easily
164 into mechanistic models of tree function. New terrestrial laser scanning methods offer a
165 way to significantly advance how we scale tree physiology from limb to tree, by enabling the
166 rapid high-resolution measurement of full tree structure, for hundreds of trees (Meir et al.
167 2017). Laser-scanning methods are advancing rapidly and will likely deliver tree structural
168 data for large forested areas within a few years, with wide application in ecological and
169 forestry science (Disney ~~et al.~~ 2018).

170
171 Moving up further-still in scale, from tree to ecosystem, the integration of such a
172 comprehensive description of tree form with the new mechanistic detail that is embedded
173 in TReSpire should help in specifying the carbon cost and climate-responsiveness of woody
174 tissue respiration at large scale. These kinds of new datasets are urgently needed to
175 examine how land-surface models perform under climate change (*cf.* Huntingford et al.
176 2017). Until now such vegetation models have made the weakly-tested assumption that the
177 respiration of woody tissue can be derived simply from leaf respiration rates, based on
178 biomass, tissue nitrogen concentration and temperature. TReSpire is a new addition to the
179 toolbox needed to determine these key components of the carbon budget of trees and
180 forests, and thereby advance how we understand the fundamental respiration cost of tree
181 size, and how this bears upon land-atmosphere interactions.

182

183 Acknowledgements

184 The authors gratefully acknowledge funding support from: ARC DP170104091~~DP17XXX~~, and
185 NERC-UK NE/L007924/1 (PM); MINECO-Spain, CGL2013-46808-R and CGL2017-89149-C2-1-
186 R XXX(MM); and an ANU doctoral scholarship (SIC).

187

Formatted: Font: 12 pt

Formatted: Font: 12 pt

188

189

190 **References**

191 Amthor J. (2000). The McCree–de Wit–Penning de Vries–Thornley respiration paradigms: 30
192 years later. *Annals of Botany* 86: 1–20.

193 Angert A, Muhr J, Juarez RN, Muñoz WA, Kraemer G, Santillan JR et al. (2012). Internal
194 respiration of Amazon tree stems greatly exceeds external CO₂ efflux. *Biogeosciences* 9,
195 4979–4991.

196 Campioli M, Malhi Y, Vicca S, Luysaert S, Papale D, Penuelas J et al. (2016) Evaluating the
197 convergence between eddy-covariance and biometric methods for assessing carbon budgets
198 of forests. *Nature Communications* 7: 13717.

199 Cannell MGR, Thornley JHM. 2000. Modelling the Components of Plant Respiration: Some
200 Guiding Principles. *Annals of Botany* 85: 45–54.

201 Chambers JQ, Tribuzy ES, Toledo LC, Crispim BF, Higuchi N, Dos Santos J, Araújo AC, Kruijt B,
202 Nobre AD, Trumbore SE (2004). Respiration from a tropical forest ecosystem: partitioning of
203 sources and low carbon use efficiency. *Ecological Applications* 14: 72–88.

204 Cuny HE, Rathgeber CBK, Frank D, P Fonti, H Mäkinen, P Prislan et al. (2015) Woody biomass
205 production lags stem-girth increase by over one month in coniferous forests. *Nature Plants*
206 1: 15175.

207 De Schepper V, Steppe K (2010). Development and verification of a water and sugar
208 transport model using measured stem diameter variations. *Journal of Experimental Botany*,
209 61, 2083–2099. DOI: 10.1093/jxb/erq018

210 Disney M (2019). Terrestrial Li DAR: a three-dimensional revolution in how we look at trees.
211 *New Phytologist* 222 (4), 1736–1741. DOI: 10.1163/22941932-90000161.

212 Fan, HL, McGuire, MA, Teskey, RO (2017). Effects of stem size on stem respiration and its
213 flux components in yellow-poplar (*Liriodendron tulipifera* L.) trees. *Tree Physiology*, 37,
214 1536–1545. DOI: 10.1093/treephys/tpx084.

215 Hilman B, Muhr J, Trumbore SE, Carbone MS, Yuval P, Joseph S. 2018. Comparison of CO₂
216 and O₂ fluxes demonstrate retention of respired CO₂ in tree stems from a range of tree
217 species. *Biogeosciences* 1: 1–30.

218 Hölttä T, Vesala T, Sevanto S, Perämäki M, Nikinmaa E (2006) Modeling xylem and phloem
219 water flows in trees according to cohesion theory and Münch hypothesis. *Trees - Structure*
220 *and Function* 20: 67–78.

221 Hölttä T, Mäkinen H, Nojd P, Makela A, Nikinmaa E (2010). A physiological model of
222 softwood cambial growth. *Tree Physiology*, 30, 1235–1252. DOI: 10.1093/treephys/tpq068

223 [Huntingford C, Atkin OK, Martinez-de la Torre A, Mercado LM, Heskell, MA et al. \(2017\).
224 Implications of improved representations of plant respiration in a changing climate. *Nature*
225 *Communications*, 8:1602. DOI: 10.1038/s41467-017-01774-z](#)

226 Levy, PE, Jarvis, PG (1998). Stem CO₂ fluxes in two Sahelian shrub species (*Guiera*
 227 *senegalensis* and *Combretum micranthum*). *Functional Ecology*, 12, 107–116. DOI:
 228 10.1046/j.1365-2435.1998.00156.x

229 Levy PE, Meir P, Allen SJ, Jarvis PG (1999) The effect of aqueous transport of CO₂ in xylem
 230 sap on gas exchange in woody plants. *Tree Physiology* 19: 53–59.

231 Meir P & Grace J (2002). Scaling relationships for woody tissue respiration in two tropical
 232 rain forests. *Plant Cell and Environment*, 25, 963–973.

233 Meir P, Shenkin A, Disney M, Rowland L, Malhi Y, Herold M, da Costa ACL. 2017. Plant
 234 structure-function relationships and woody tissue respiration: Upscaling to forests from
 235 laser-derived measurements. In: Tcherkez G, Ghashghaie J, eds. *Plant Respiration: Metabolic*
 236 *fluxes and carbon balance*. Cham: Springer International Publishing, 89–105.

237 Mencuccini M, Holttä T, Sevanto S, Nikinmaa E (2013). Concurrent measurements of change
 238 in the bark and xylem diameters of trees reveal a phloem-generated turgor signal. *New*
 239 *Phytologist*, 198, 1143–1154. DOI: 10.1111/nph.12224

240 Mencuccini M, Salmon Y, Mitchell P, Hölttä T, Choat B, Meir P, O’Grady A, Tissue D, Zweifel
 241 R, Sevanto S, et al. 2017. An empirical method that separates irreversible stem radial growth
 242 from bark water content changes in trees: theory and case studies. *Plant, Cell &*
 243 *Environment* 40: 290–303.

244 Metcalfe DB, Meir P, Aragao L, Lobo-do-Vale R, Galbraith D, Fisher RA et al. (2010) Shifts in
 245 plant respiration and carbon use efficiency at a large-scale drought experiment in the
 246 eastern Amazon. *New Phytologist*, 187, 608–621.

247 Mori S, Yamaji K, Ishida A, Prokushkin SG, Masyagina OV, Hagihara A et al. (2010) Mixed-
 248 power scaling of whole-plant respiration from seedlings to giant trees. *Proceedings of the*
 249 *National Academy of Sciences USA*, 107, 1447–1451.

250 O’Leary BM, Asao S, Millar AH, Atkin OK (2019). Core principles which explain variation in
 251 respiration across biological scales. *New Phytologist*, 222, 670–686. DOI:
 252 10.1111/nph.15576

253 Reich PB, Tjoelker MG, Machado JL, Oleksyn J (2006) Universal scaling of respiratory
 254 metabolism, size and nitrogen in plants. *Nature* 439: 457 —461.

255 Rossi S, Deslaurier A, Anfodillo T (2006). Assessment of cambial activity and xylogenesis by
 256 microsampling tree species: An example at the alpine timberline. *IAWA Journal*, 27, 383–
 257 394.

258 Ryan MG, Lavigne MB, Gower ST (1997) Annual carbon balance cost of autotrophic
 259 respiration in boreal forest ecosystems in relation to species and climate. *Journal of*
 260 *Geophysical Research* 102: 28871–28883.

261 Steppe K, Saveyn A, McGuire MA, Lemeur R, Teskey RO. 2007. Resistance to radial CO₂
 262 diffusion contributes to between-tree variation in CO₂ efflux of *Populus deltoides* stems.
 263 *Functional Plant Biology* 34: 785–792.

264 Teskey RO, Saveyn A, Steppe K, McGuire MA. 2008. Origin, fate and significance of CO₂ in
 265 tree stems. *New Phytologist* 177: 17–32.

266 Thornley JHM (1970) Respiration, growth, and maintenance in plants. *Nature* 227, 304–305.
267 [West PW \(2019\). Do increasing respiratory costs explain the decline with age of forest](#)
268 [growth rate? *Journal of Forestry Research*. DOI: 10.1007/s11676-019-01020-w](#)
269 Yang J, He Y, Aubrey DP, Zhuang Q, Teskey RO. 2016. Global patterns and predictors of stem
270 CO₂ efflux in forest ecosystems. *Global Change Biology* 22: 1433–1444.
271 ~~Yoneda T (1993) Surface area of woody organs of an evergreen broadleaf forest tree in~~
272 ~~Japan *Southeast Asian Journal of Plant Research* 106: 229–237.~~
273 Zweifel R. (2016) Radial stem variations—a source of tree physiological information not fully
274 exploited yet. *Plant, Cell & Environment* 39, 231–232.